

1. Achieving Land Degradation Neutrality through Soil-centred Restoration and Management Strategies

Dr. Yashika Mandela¹, Dr. Uday Sharma¹, Dr. Anusuiya Panda¹, Dr. Shradha Chauhan¹, Dr. Kiran Masta¹

¹Ph.D. Scholar, Division of Soil Science and Water Management, Dr Yashwant Singh Parmar University of Horticulture and Forestry, Nauni (Solan), Himachal Pradesh

1. Introduction

Land degradation, defined by the United Nations Convention to Combat Desertification (UNCCD) as the reduction or loss of the biological or economic productivity and complexity of rainfed cropland, irrigated cropland, or rangeland, pasture, forest, and woodlands, represents one of the most pressing global environmental and socio-economic crises of the 21st century. Currently, degradation affects over 25% of the Earth's total land area, directly impacting the livelihoods of billions of people, predominantly in developing economies. The resulting annual economic cost, attributable to the loss of ecosystem services, is estimated at hundreds of billions of U.S. dollars. This pervasive challenge undermines global efforts to achieve food security, exacerbates poverty, contributes significantly to climate change by releasing soil carbon, and accelerates biodiversity loss.

The gravity of this global issue led to the establishment of Land Degradation

Neutrality (LDN) as a specific target, Target 15.3, within the United Nations Sustainable Development Goals (SDGs). LDN is conceptually defined as a state in which the amount of healthy, productive land resources necessary to support ecosystem services and food security remains stable or increases within specified spatial and temporal scales (UNCCD, 2017). This aspirational target requires a fundamental shift in land management philosophy, moving from a purely extractive model to one based on sustainability and restoration. LDN is essentially a spatial balancing act: it seeks to balance anticipated losses from degradation (hotspots) with planned gains from restoration and sustainable management (bright spots), aiming for a net-zero or net-positive change in the health of land resources.

Achieving this complex state of neutrality necessitates a strategic, targeted approach. This article posits that the successful implementation of the LDN agenda

depends on placing soil health at the core of all policy and intervention strategies. The soil, a finite, non-renewable resource over human timescales, is the foundation of terrestrial ecosystems. It provides crucial ecosystem services, including nutrient cycling, water storage, and acts as the most significant terrestrial carbon sink. When soil health declines, as evidenced by erosion, compaction, nutrient depletion, and loss of Soil Organic Carbon (SOC), the land system's functional capacity collapses, leading directly to land degradation.

Therefore, the LDN imperative translates into an SOC imperative and a soil functional imperative. The strategy must be two-fold:

- a. Prevention:** Proactively protecting currently productive soils from further degradation.
- b. Restoration:** Actively rebuilding the productivity of already degraded soils by focusing on the enhancement of critical soil properties (physical, chemical, and biological).

1. The Centrality of Soil in Land Degradation and Restoration

Soil, often called the skin of the Earth, is a complex and finite natural resource critical to sustaining human life and terrestrial ecosystems. Its functionality is intrinsically linked to its ability to provide essential

ecosystem services, including biomass production, nutrient cycling, water purification and storage, and climate regulation through carbon sequestration. The process of land degradation fundamentally reflects the deterioration of soil health and its functional capacity, which is typically slow to reverse and highly costly to repair.

1.1. Manifestations and Mechanisms of Soil Degradation

Land degradation is a multifaceted process manifesting through the decline of three key soil property domains:

- a. Physical Degradation:** This primarily involves soil erosion and compaction. Erosion, driven by water and wind, selectively removes the fertile topsoil enriched with fine particles, clay minerals, and organic matter. This leads to reduced rooting depth and overall water-holding capacity. Compaction, often caused by heavy machinery, decreases soil porosity and increases bulk density, severely restricting water infiltration and root penetration, thereby increasing runoff and increasing the risk of erosion.
- b. Chemical Degradation:** This includes nutrient depletion, where the concentration of essential

elements (N, P, K, S) and micronutrients (Zn, Mn, Cu, Fe) falls below the critical level required for optimal plant growth. Equally damaging are processes such as salinisation (an increase in soluble salts, indicated by a high Electrical Conductivity (EC), sodification (excess sodium), and acidification (a drop in pH). These chemical imbalances disrupt nutrient availability, creating toxic environments that stress soil organisms and cultivated crops.

- c. **Biological Degradation:** The loss of Soil Organic Carbon (SOC) is the single most critical biological indicator of degradation. Intensive tillage accelerates the oxidation of SOC into CO₂, depleting the soil's organic matter pool. This decline cripples the soil's structure, diminishes its capacity to hold water and nutrients, and reduces the diversity and activity of the soil microbiome. A reduction in microbial biomass and functional diversity impairs decomposition and nutrient cycling processes.

1.2. Restoring Soil Health as the LDN Imperative

Restoring degraded land, therefore, is synonymous with rebuilding soil health by

targeting these three domains. A high and stable SOC content, a resilient aggregate structure, a balanced pH and EC, and a robust biological community characterise a healthy soil.

Key Soil Properties and the LDN Framework:

The commitment to LDN requires measurable metrics. The UNCCD's LDN monitoring framework mandates explicitly the use of Soil Organic Carbon (SOC) Stock as one of its three core global indicators. SOC is prioritised because:

- a. It is directly linked to soil fertility (chemical and biological health).
- b. It acts as a primary binding agent for soil aggregates (physical health).
- c. It serves as a massive terrestrial carbon sink, linking LDN directly to climate change mitigation (SDG 13).

Advanced techniques, such as soil spectroscopy and remote sensing, are becoming indispensable for rapidly and cost-effectively measuring a suite of critical soil properties (including SOC, EC, pH, and available nutrients like P, K, S, Zn, Mn, Cu, and Fe), enabling targeted, precision restoration interventions at scale. By strategically focusing on practices that enhance SOC and improve the soil's capacity to regulate water and nutrients, we move toward achieving a net-positive

balance in land health, the foundational objective of LDN.

2. Soil-Centred Restoration and Management Strategies for LDN

Achieving LDN requires the widespread adoption of Sustainable Land Management (SLM) practices that directly address the drivers of soil degradation. These practices focus on minimising soil disturbance, maximising organic matter return, and optimising the efficient use of water and nutrients. Three cornerstone soil-centred strategies are detailed below.

2.1. Conservation Agriculture (CA)

Conservation Agriculture is a transformative farming system defined by three interconnected principles that collectively enhance the physical, chemical, and biological dimensions of soil health:

- a. Minimum Soil Disturbance (No-Till/Reduced Tillage):** This practice is key to maintaining soil aggregate stability and minimising the rapid oxidation and loss of Soil Organic Carbon (SOC). By avoiding mechanical shearing of soil aggregates, no-till preserves macropores, which are vital for enhancing water infiltration rates and minimising runoff, thereby acting as a primary erosion control measure.

- b. Permanent Soil Cover:**

Maintaining a continuous layer of crop residues or living cover crops (mulch) protects the soil from the kinetic energy of rainfall, which causes dispersion and crusting. This cover significantly conserves soil moisture, moderates soil temperature, and provides a continuous surface input of organic material, which is essential for feeding soil biota and supporting SOC accumulation.

- c. Crop Rotation/Diversification:**

Varied rooting patterns and nutrient demands from diverse crops help alleviate compaction and improve the distribution of nutrients throughout the soil profile. The incorporation of legumes into the rotation is critical for biological nitrogen fixation (BNF), reducing the need for high-energy nitrogen fertilisers and improving overall nutrient balance (e.g., N content).

The adoption of CA leads to demonstrable improvements in soil structure, increased SOC stocks, and greater water-use efficiency, directly countering the physical and biological




Figure 1: Conceptual Model of Soil Health Improvement under Conservation Agriculture

degradation aspects of the LDN challenge.

Table 1: Key Principles and Benefits of Conservation Agriculture in the Context of LDN

Principle	Impact on Soil Health	LDN Relevance
Minimum Tillage	Preserves soil structure; reduces SOC loss via oxidation.	Prevents degradation; maintains SOC stock (LDN Indicator).
Permanent Soil Cover	Controls erosion, conserves moisture, and increases organic matter.	Reverses land productivity decline (LDN Indicator).
Crop Diversification	Improves nutrient balance, enhances soil biodiversity, and reduces disease pressure.	Enhances ecosystem resilience; sustains long-term productivity.

2.2. Agroforestry Systems

Agroforestry, the intentional integration of trees and shrubs with crops and/or livestock, represents a potent bio-engineering solution for soil restoration and stabilisation, particularly in marginal or degraded lands.

a. Soil Stabilisation and Hydrology:

Deep tree root systems provide mechanical reinforcement to the soil profile, dramatically reducing the risk of mass movement and stabilising soil against deep erosion.

The presence of trees alters the microclimate and improves the overall hydrological cycle, leading to greater SOC accumulation and enhanced water use efficiency compared to treeless monocultures.

b. Nutrient Cycling and Acquisition:

Trees function as natural nutrient

pumps, accessing nutrients and water from deeper soil horizons unavailable to shallow-rooted annual crops. These nutrients are brought to the surface through litterfall and decomposition, thereby improving the availability of macro- and micronutrients (such as P and K) in the topsoil. Furthermore, leguminous trees contribute significantly to the soil's N capital through symbiotic BNF.

By fostering greater spatial and temporal biomass diversity, agroforestry systems are highly effective at reversing erosion and substantially increasing overall SOC sequestration potential.

2.3. Integrated Nutrient Management (INM)

INM is a holistic strategy that seeks to sustain crop productivity while protecting the environment by optimising the combined use of all nutrient sources.

- a. SOC and Soil Structure Enhancement:** The rigorous use of organic resources (farmyard manure, compost, green manures, and biochar) directly contributes stable Carbon to the soil matrix. This input is crucial for forming stable soil aggregates,

improving soil tilth and water-holding capacity, and reducing bulk density.

- b. Precision and Efficiency:** INM advocates for site-specific nutrient management (SSNM), which is essential for correcting specific chemical degradation issues. Using diagnostics such as soil testing for pH and EC allows for targeted application of lime or gypsum to address acidification or salinisation, respectively. The principle is to apply the right amount of inorganic fertiliser to supplement organic sources, thereby maximising Nutrient Use Efficiency (NUE) and minimising environmental impacts from nutrient leaching.

Table 2: Mechanisms of Soil-centred Strategies in Reversing Degradation

Strategy	Primary Degradation Addressed	Key Mechanism for Restoration	Measurable Soil Improvement
Agroforestry	Erosion, Biological, Chemical	Deep rooting, N-fixation, continuous litterfall.	SOC increased, and P and K availability improved.
Conservation Agriculture	Physical (Erosion/Compaction), Biological	Reduced tillage, permanent cover, residue incorporation.	Aggregate stability, increased water infiltration, and higher SOC.
INM	Chemical (Nutrient)	Balanced use of organic	Optimal pH and

	depletion, pH imbalance)	and inorganic inputs; SSNM.	EC; sustained levels of P, K, S.
--	--------------------------	-----------------------------	----------------------------------

3. Policy and Implementation Frameworks for LDN

The technical feasibility of soil-centred restoration strategies must be complemented by robust policy and institutional frameworks to ensure scalable and sustained adoption. Achieving LDN is not merely a technical challenge but a complex socio-economic and governance task that requires alignment across multiple governmental sectors.

3.1. Governance and National LDN Alignment

The commitment to LDN originates at the global level but is realised through decisive national action. Countries that are signatories to the UNCCD are mandated to formulate National LDN Targets, which are voluntary but critically important for establishing baselines and setting specific, time-bound goals. These targets compel governments to:

- a. Mainstream SLM:** Integrate soil health objectives into broader national development plans, including those for agriculture, food security, climate change adaptation, and biodiversity conservation.
- b. Legal and Institutional Reforms:** Review and reform land tenure laws to

provide long-term security to land users, as insecure tenure is a significant disincentive for investment in soil-restorative practices like agroforestry, which require long time horizons.

c. Cross-Sectoral Coordination:

Establish formal mechanisms to ensure collaboration between ministries (Agriculture, Environment, Finance) to prevent policy contradictions and maximise the co-benefits of LDN efforts.

3.2. Economic Incentives and Sustainable Financing

The transition to soil-restorative practices like Conservation Agriculture (CA) involves initial costs and a potential lag before full economic benefits are realised, necessitating supportive financial policy. Effective mechanisms include:

- a. Direct Financial Support:** Offering subsidies for procuring specialised equipment (e.g., no-till seeders) and inputs (e.g., cover crop seeds). This lowers the initial barrier to adoption for smallholder farmers.
- b. Payment for Ecosystem Services (PES):** Developing schemes where public or private entities compensate farmers for the environmental services derived from their soil management. The emergent Voluntary Carbon Market (VCM) offers a significant financial opportunity, providing farmers with revenue based on

verifiable increases in Soil Organic Carbon (SOC) stocks —a direct link between soil health and climate finance.

c. Risk Mitigation and Credit Access: Providing contingency planning and index-based crop insurance that rewards climate-resilient practices. Furthermore, policies must facilitate access to affordable credit for rural populations to finance the upfront capital required for SLM investments.

3.3. Data-Driven Monitoring and Adaptive Management

Robust monitoring is non-negotiable for LDN accountability and effective policy adaptation. The UNCCD framework relies on tracking the three key indicators: land cover, land productivity, and SOC stock.

- **Leveraging Spectral Technology:** The user's area of research, the integration of remote sensing and soil spectroscopy, is paramount for this component. Remote sensing provides wide-area spatial monitoring of land cover change and productivity trends. Crucially, Proximal and Mid-Infrared (NIR-MIR) Soil Spectroscopy, combined with advanced machine learning, offers a rapid, low-cost, and non-destructive method for routinely measuring essential soil properties (e.g., SOC, EC, pH, P, K, S, Zn, Mn, Cu, Fe) at a high spatial density. This allows for:

a. Accurate Baseline Establishment: Creating precise maps of the current soil health status.

b. Targeted Intervention: Identifying specific "hotspots" of degradation requiring immediate attention.

c. Verification and Reporting: Providing verifiable data for national SOC inventory reporting and supporting participation in carbon markets.

This data-driven approach enables adaptive governance, where policies and on-the-ground strategies are continuously adjusted based on scientific evidence of what works and where degradation pressures persist. By institutionalising this feedback loop, the LDN process remains dynamic and effective in a changing environment.

Table 3: Policy and Technological Tools for LDN Implementation and Monitoring

Category	Tool/Mechanism	LDN Function	Impact on Soil-Centric Goals
Policy	National LDN Targets (SDG 15.3)	Provides the political mandate and structure for action.	Focuses national resources on preserving SOC and land productivity.
Finance	Payment for Ecosystem Services (PES)	Incentivises the voluntary adoption of SLM practices by farmers.	Compensates for the long-term investment required for SOC

			restoration.
Technology	Remote Sensing (Satellite Imagery)	Monitors large-scale land cover and productivity trends.	Tracks the extent of healthy land resources (LDN Indicator 1 & 2).
Technology	Soil Spectroscopy and Machine Learning	Provides rapid, accurate, and cost-effective measurement of soil properties.	Verifies changes in SOC, pH, EC, P, and K for LDN reporting and carbon markets.

5. Conclusion

Achieving Land Degradation Neutrality (LDN) is not merely a land-use policy objective; it is an imperative for global sustainability, food security, and climate resilience. As demonstrated, the success of the LDN framework is intrinsically tied to the successful implementation of soil-centred restoration and management strategies. By focusing on the fundamental health of the soil — specifically through the enhancement of Soil Organic Carbon (SOC) stocks, soil structure stability, and biological diversity — we directly address the physical, chemical, and biological drivers of degradation.

The wide-scale adoption of proven Sustainable Land Management (SLM) practices, including Conservation Agriculture (CA), Agroforestry, and Integrated Nutrient Management (INM),

offers a viable and scalable pathway. These systems work synergistically to minimise soil loss, maximise water use efficiency, and sequester atmospheric carbon, thereby delivering crucial co-benefits beyond LDN, such as climate change mitigation and enhanced biodiversity.

However, realising LDN requires moving beyond technical solutions. It demands strong policy frameworks that translate global targets into actionable national plans, supported by financial mechanisms (such as PES and carbon markets) that incentivise farmers and land managers to shift towards soil-friendly practices. Crucially, the future of LDN monitoring and reporting will depend on advanced technologies, such as integrating remote sensing and spectral data to accurately, rapidly, and cost-effectively estimate key soil health indicators (SOC, pH, EC, etc.).

References

- Lal, R. (2015). Soil carbon sequestration and land degradation neutrality. *Carbon Management*, 6(3), 119–125.
- Minasny, B., McBratney, A. B., Bell, M. J., & Roudier, P. (2024). Spectroscopy and machine learning: Revolutionising soil quality monitoring for sustainable resource management. In P. K. Pande (Ed.), *Natural Resource Monitoring, Planning and Management Based on*

- Advanced Programming* (pp. 199–223). Springer.
- Shi, T., Liu, Y., Wu, G., Wang, Y., & Zuo, H. (2023). Using machine-learning algorithms to predict soil organic carbon content from combined remote sensing imagery and laboratory Vis-NIR spectral datasets. *Remote Sensing*, 15(17), 4264.
- Taghizadeh-Mehrjardi, R., Minasny, B., & Ghadiri, H. (2023). Spatial prediction of physical and chemical properties of soil using optical satellite imagery: A state-of-the-art hybridisation of deep learning algorithms. *Frontiers in Environmental Science*, 11.
- UNCCD (United Nations Convention to Combat Desertification). (2017). *Scientific Conceptual Framework for Land Degradation Neutrality*. A report of the Science-Policy Interface. UNCCD.
- Venter, Z. S., Hengl, T., & van Rijn, I. (2021). Digital soil mapping of soil organic carbon in agricultural areas using remote sensing and machine learning. *Geoderma*, 391, 114959.