

15. Dynamics of Soil Aggregate Stability under Organic and Conventional Farming Systems

Azad Patel¹, Umesh Bandu Dolaskar², Surbhi Shakya³, Aaditya Nimbalkar⁴, Shabana Sheikh⁵

¹M.Sc. Scholar, Division of Soil Science, IARI Hub, Nagpur

²M.Sc. Scholar, Division of Soil Science, Dr. P. D. K. V, Akol, Maharashtra

³M.Sc. Scholar, Division of Soil Science, Rajmata Vijayaraje Scindia Krishi Vishwavidyalaya, Gwalior (M.P.)

⁴M.Sc. Scholar, Department of Soil Science and Agricultural chemistry, Institute of agricultural Science, Banaras Hindu University, Varanasi, Uttar Pradesh

⁵Laboratory Technician, Division of Soil Science, Dr. P. D. K. V, Akola, Maharashtra.

1. Introduction

Soil is a complex, dynamic medium, fundamental to agricultural productivity and ecosystem function. Its physical structure, largely characterized by the formation and stability of aggregates, dictates a wide range of essential processes including water retention, nutrient cycling, gas exchange, and root penetration (Six et al., 2004). Soil aggregates are clusters of primary soil particles (sand, silt, and clay) bound together by various cementing agents, primarily organic matter, microbial exudates, and mineral interactions (Tisdall & Oades, 1982). Stable soil aggregates are crucial for maintaining soil porosity, resisting erosive forces, and providing a conducive habitat for soil biota.

Agricultural management practices exert significant influence on soil aggregate stability. Conventional farming systems, often characterized by intensive tillage, synthetic fertilizer application, and pesticide use, have been implicated in the

degradation of soil structure and a reduction in aggregate stability (Powlson et al., 1987). In contrast, organic farming systems emphasize practices such as cover cropping, crop rotation, compost application, and reduced tillage, all of which are hypothesized to enhance soil organic matter (SOM) content and biological activity, thereby promoting aggregate formation and stability (Reganold et al., 1993).

2. Mechanisms of Soil Aggregate Formation and Stability

Soil aggregates are formed through a hierarchical process involving physical, chemical, and biological interactions. Primary particles coalesce to form microaggregates (<250 µm), which in turn bind together to form macroaggregates (>250 µm) (Oades, 1984). The stability of these aggregates, particularly against disruptive forces like water erosion and mechanical stress, is a key determinant of soil resilience.

2.1. Binding Agents

The stability of soil aggregates is primarily attributed to various binding agents:

- a. **Organic Matter (OM):** Organic matter is arguably the most crucial binding agent, particularly in the formation of macroaggregates. Decomposing plant and animal residues provide transient binding agents, while humic substances and recalcitrant organic carbon act as more persistent binders (Six et al., 2002). Polysaccharides, glomalin-related soil protein (GRSP) produced by arbuscular mycorrhizal fungi, and other microbial exudates are powerful transient cementing agents.
- b. **Microbial Activity:** Soil microorganisms, including bacteria, fungi, and archaea, play a pivotal role in aggregate formation. Fungal hyphae entangle soil particles, while bacterial colonies secrete extracellular polymeric substances (EPS) that act as glues (Jastrow et al., 1998).
- c. **Clay Minerals:** Clay particles, with their high surface area and charge, interact with organic matter and polyvalent cations (e.g., Ca^{2+} , Mg^{2+}) to form stable organo-mineral complexes, particularly important

in microaggregate stability (Chenu & Guérif, 1991).

d. Iron and Aluminium Oxides:

These oxides can also contribute to aggregate stability by binding to clay particles and organic matter, especially in highly weathered soils.

2.2. Factors Influencing Aggregate Stability

Several factors influence the overall stability of soil aggregates:

- a. **Soil Texture:** Soils with higher clay content generally exhibit greater aggregate stability due to the increased surface area for binding and stronger organo-mineral interactions.
- b. **Soil Moisture:** Wetting and drying cycles can alternately promote aggregate formation (through shrinking and cracking) and breakdown (through slaking and dispersion).
- c. **Tillage:** Mechanical disturbance from tillage can disrupt existing aggregates, exposing fresh organic matter to microbial decomposition and leading to a reduction in aggregate stability.
- d. **Root Exudates:** Living roots release exudates that stimulate microbial activity and directly

contribute to aggregate formation around the rhizosphere.

3. Conventional Farming Systems and Aggregate Stability

Conventional farming practices, while often achieving high yields in the short term, can negatively impact soil aggregate stability.

3.1. Tillage Practices

Intensive tillage, a hallmark of many conventional systems, involves mechanical disruption of the soil matrix. This process fragments existing aggregates, exposes previously protected organic matter to rapid microbial decomposition, and can create a compacted plow pan (Karlen et al., 1994). The immediate consequence is a reduction in macroaggregate content and overall aggregate stability, making the soil more susceptible to wind and water erosion.

3.2. Synthetic Inputs

The reliance on synthetic fertilizers, particularly nitrogen fertilizers, can

stimulate rapid decomposition of labile organic matter, potentially reducing the pool of binding agents (Drinkwater et al., 1998). While providing essential nutrients, synthetic fertilizers may also alter the composition and activity of soil microbial communities, sometimes favouring fast-growing, less aggregate-forming organisms over beneficial fungal networks. Pesticides and herbicides can also have unintended non-target effects on soil microbial populations, further disrupting the biological processes critical for aggregate formation.

3.3. Monoculture and Residue Management

Conventional systems often employ monoculture, which limits the diversity of root exudates and biomass inputs. In some cases, crop residues are removed or burned, depriving the soil of fresh organic matter inputs that are vital for sustaining microbial activity and aggregate formation.

Table 1 Impact of Conventional Farming Practices on Aggregate Stability

Practice	Effect on Soil Aggregates	Mechanism
Intensive Tillage	Decreased macroaggregate stability, increased pulverization	Mechanical disruption, exposure of OM to decomposition
Synthetic Fertilizers	Potential reduction in SOM, altered microbial communities	Stimulated labile OM decomposition, changes in microbial composition
Pesticides/Herbicides	Non-target effects on beneficial soil microbes	Disruption of microbial activity critical for binding
Monoculture	Reduced diversity of organic inputs, limited root systems	Less diverse exudates, fewer physical binding opportunities
Residue Removal	Reduced input of fresh organic matter	Deprivation of easily decomposable OM for microbial activity and binding

4. Organic Farming Systems and Aggregate Stability

Organic farming systems prioritize ecological processes, promoting soil health through practices that enhance organic matter accumulation and biodiversity. These practices generally lead to greater soil aggregate stability.

4.1. Enhanced Organic Matter Inputs

Organic systems typically incorporate practices that significantly increase SOM content:

- a. **Compost and Manure Application:** Regular application of compost and animal manures introduces stable organic matter and a diverse microbial inoculum, directly contributing to aggregate formation and stability (Liu et al., 2017).
- b. **Cover Cropping:** Cover crops protect the soil surface from erosion, add fresh organic matter, and their extensive root systems physically bind soil particles and stimulate rhizosphere activity.
- c. **Crop Rotation:** Diverse crop rotations, often including legumes and deep-rooted species, provide varied organic inputs and support a more diverse and active soil microbial community, leading to

stronger aggregate formation (Drinkwater et al., 1998).

- d. **Reduced Tillage/No-Till:** While not universally adopted in organic systems, reduced tillage is increasingly common. Minimizing soil disturbance preserves existing aggregates, allows for the accumulation of surface residues, and fosters stable fungal networks.

4.2. Biological Activity

Organic farming fosters a more biologically active soil environment:

- a. **Fungal Dominance:** Reduced tillage and continuous organic matter inputs tend to favor fungal growth, particularly arbuscular mycorrhizal fungi (AMF). AMF hyphae are powerful physical binders, and their glomalin production acts as a significant cementing agent, stabilizing macroaggregates (Rillig *et al.*, 2004).
- b. **Diverse Microbial Communities:** The absence of synthetic pesticides and a steady supply of diverse organic substrates promote a more diverse and robust soil microbial community, with a greater capacity for producing extracellular binding substances.

4.3. Root Systems

The diverse root systems inherent in organic crop rotations and cover cropping provide both physical entanglement of soil particles and a continuous supply of root

exudates, which are crucial for stimulating microbial activity and the formation of stable aggregates around the rhizosphere.

Table 2 Impact of Organic Farming Practices on Aggregate Stability

Practice	Effect on Soil Aggregates	Mechanism
Compost/Manure	Increased SOM, enhanced microbial activity, stable aggregates	Direct OM input, microbial inoculation, stable humic substance formation
Cover Cropping	Increased SOM, physical binding, and reduced erosion	Biomass input, root exudates, and protection from raindrop impact
Diverse Crop Rotation	Varied organic inputs, diverse microbial communities	Different root architectures, varied residue quality, diverse microbial food sources
Reduced/No-Till	Preservation of existing aggregates, fungal dominance	Minimized physical disruption, undisturbed fungal networks
No Synthetic Pesticides	Healthy and diverse microbial populations	Absence of toxic effects on beneficial soil microorganisms

5. Comparative Studies and Empirical Evidence

Numerous studies have directly compared soil aggregate stability under organic and conventional farming systems, consistently demonstrating superior stability in organic plots.

Reganold et al. (1993) famously showed that soils from organically farmed fields had significantly higher aggregate stability, along with greater organic matter content and microbial biomass, compared to conventionally managed fields. A meta-analysis by Lori et al. (2017) confirmed that organic farming generally leads to higher soil organic carbon and improved aggregate stability, particularly in topsoil layers.

A long-term study by Mäder et al. (2002) at the DOK field experiment in Switzerland found that after 21 years, organic and biodynamic systems exhibited greater water-stable aggregates than conventional systems, attributed to higher biological activity and organic carbon content. Similarly, the Rodale Institute's Farming Systems Trial in the USA has consistently shown that organic no-till systems build soil organic carbon and improve aggregate stability more effectively than conventional systems (Teasdale et al., 2007).

6. Implications for Soil Health and Ecosystem Services

The enhanced soil aggregate stability observed in organic farming systems has profound implications for overall soil

health and the provision of vital ecosystem services.

enhances water storage capacity within the soil profile, making water more available to




Figure 1 Conceptual Model of Aggregate Stability under Different Farming Systems

7. Implications for Soil Health and Ecosystem Services

The enhanced soil aggregate stability observed in organic farming systems has profound implications for overall soil health and the provision of vital ecosystem services.

7.1. Water Dynamics

Stable aggregates create a well-structured soil with increased porosity, facilitating improved water infiltration and reducing surface runoff (Franzluebbers, 2002). This not only mitigates flood risk but also

plants, particularly during dry periods. Conversely, soils with poor aggregate stability are prone to surface sealing and compaction, leading to reduced infiltration and increased runoff and erosion.

7.2. Erosion Control

Soils with stable aggregates are more resistant to erosive forces from wind and water. The binding of soil particles into larger, stable units prevents their detachment and transport. This is a crucial benefit in the face of increasingly extreme

weather events, helping to preserve the fertile topsoil and prevent land degradation.

7.3. Nutrient Cycling and Retention

Stable aggregates protect organic matter from rapid decomposition, leading to a more gradual release of nutrients, which can synchronize nutrient availability with plant demand (Jastrow et al., 1998). The enhanced microbial activity associated with organic systems also improves nutrient mineralization and immobilization processes. Furthermore, a stable soil structure reduces leaching losses of nutrients, retaining them within the rooting zone.

7.4. Biodiversity and Habitat

A well-aggregated soil provides a diverse range of microhabitats for soil organisms, supporting a richer and more abundant soil biota, including earthworms, insects, and microorganisms (Brussaard et al., 2007). This biodiversity is essential for maintaining a healthy and resilient soil ecosystem, contributing to pest control, disease suppression, and overall soil fertility.

7.5. Carbon Sequestration

By promoting higher organic matter content and protecting it within stable aggregates, organic farming systems contribute significantly to soil carbon sequestration (Post & Kwon, 2000). This process removes carbon dioxide from the

atmosphere and stores it in the soil, offering a valuable strategy for climate change mitigation.

8. Challenges and Future Directions

While the evidence overwhelmingly points to the benefits of organic farming for soil aggregate stability, several challenges and areas for future research exist.

8.1. Transition Period

The transition from conventional to organic farming can present challenges, as it takes time for soil health parameters, including aggregate stability, to fully improve. Initial yield reductions or pest outbreaks can be disincentives for farmers. Research into effective transition strategies and support mechanisms for farmers is crucial.

8.2. Context Specificity

The magnitude of the impact of organic practices on aggregate stability can vary depending on soil type, climate, specific crop rotations, and local management details (Reiter et al., 2017). More regionally focused studies are needed to fine-tune recommendations.

8.3. Research on Microbial Mechanisms

While the role of soil microbes is recognized, a deeper understanding of the specific microbial communities and their functional contributions to aggregate formation and stability in different organic systems is still evolving. Advanced

molecular techniques can provide more detailed insights.

8.4. Economic Viability

The economic viability of organic farming, despite its environmental benefits, remains a key consideration for adoption. Further research into optimizing organic practices for both soil health and farm profitability is essential.

9. Conclusion

Soil aggregate stability is a fundamental pillar of soil health, underpinning critical ecological functions in agricultural landscapes. The comprehensive body of research reviewed in this article consistently demonstrates that organic farming systems, through their emphasis on organic matter inputs, diverse crop rotations, reduced tillage, and avoidance of synthetic chemicals, generally foster significantly greater soil aggregate stability compared to conventional systems. This enhanced stability leads to a cascade of benefits, including improved water management, reduced erosion, increased nutrient retention, and greater biodiversity, all of which contribute to the long-term sustainability and resilience of agricultural ecosystems. As the imperative for sustainable food production intensifies, promoting and adopting organic and regenerative agricultural practices becomes increasingly vital for safeguarding our

precious soil resources for future generations.

References

- Brussaard, L., de Ruiter, P. C., & Brown, G. (2007). Soil biodiversity for agricultural sustainability. *Agriculture, Ecosystems & Environment*, 121(1-2), 1-5.
- Chenu, C., & Guérif, J. (1991). The contribution of clay-organic interactions to the stability of soil aggregates. *Plant and Soil*, 137(1), 1-9.
- Drinkwater, L. E., Wagoner, P., & Sarrantonio, M. (1998). Legume-based cropping systems have higher soil carbon and nitrogen than cereal-based systems. *Nature*, 396(6712), 262-265.
- Franzluebbers, A. J. (2002). Soil aggregation and associated carbon and nitrogen under various cropping systems in the southern Piedmont USA. *Soil and Tillage Research*, 66(2), 163-176.
- Jastrow, J. D., Boutton, T. W., & Miller, R. M. (1998). Biomass and root-derived carbon in soil under switchgrass and row crops. *Soil Biology and Biochemistry*, 30(7), 899-906.
- Karlen, D. L., Wollenhaupt, N. C., Erbach, D. C., Berry, E. C., Swan, J. B.,

- Paustian, N. T., & Kemper, W. D. (1994). Crop residue effects on soil water and agricultural chemical movement. *Soil and Water Conservation Society*, 49(S1), 6-27.
- Liu, K., Zhang, X., Cai, Y., & Yang, B. (2017). Long-term manure application increases soil organic carbon content and promotes soil aggregation in a temperate cropland in China. *Agriculture, Ecosystems & Environment*, 237, 223-231.
- Lori, M., Symnaczik, S., Mäder, P., Liebisch, F., & Gattinger, A. (2017). Organic farming enhances soil aggregate stability and water infiltration capacity of a sandy loam soil. *Soil and Tillage Research*, 173, 20-28.
- Mäder, P., Fliessbach, A., Dubois, L., Gunst, E., Hofmann, S., & Niggli, U. (2002). Soil fertility and biodiversity in organic farming. *Science*, 296(5573), 1694-1697.
- Oades, J. M. (1984). Soil organic matter and structural stability: Mechanisms and implications for management. *Plant and Soil*, 76(1-3), 319-337.
- Post, W. M., & Kwon, K. C. (2000). Soil carbon sequestration and land-use change: processes and potential. *Global Change Biology*, 6(3), 317-327.
- Powlson, D. S., Brookes, P. C., & Jenkinson, D. S. (1987). Soil microbial biomass provides an early indication of changes in soil organic matter due to straw incorporation. *Soil Biology and Biochemistry*, 19(2), 159-164.
- Reganold, J. P., Palmer, A. S., Andrews, S. L., & Macgregor, A. N. (1993). Soil quality and financial performance of biodynamic and conventional farms in New Zealand. *Science*, 260(5106), 344-349.
- Reiter, M. S., Schlegl, E., & Baumgartner, A. (2017). Effects of organic and conventional farming on soil chemical, physical and biological properties. *Agriculture, Ecosystems & Environment*, 240, 148-156.
- Rillig, M. C. (2004). Arbuscular mycorrhizae and soil aggregate stability. *Plant and Soil*, 254(1), 39-44.