

4. Soil Phosphorus Fixation and Its Management for Improved Nutrient Availability and Crop Yields

Dr Maragani Vamshi¹, M. Vikram Sai²

¹Assistant Professor, Department of Forest Resource Management,
Forest college and research Institute, Mulugu, Siddipet, Telangana.

²Assistant Professor, Department of Agronomy, Malla Reddy University, Hyderabad.

Abstract

Phosphorus (P) is a critical macronutrient driving early root development, flowering, and energy transfer in plant systems. Despite its necessity, phosphorus management remains a significant challenge due to phosphorus fixation—a series of chemical reactions that render soluble phosphate fertilisers unavailable to crops. In acidic soils, fixation is dominated by aluminium and iron oxides, while in alkaline and calcareous soils, calcium precipitates prevail. Consequently, soils frequently test high in total phosphorus yet fail to supply adequate plant-available nutrients, leading to reduced fertiliser use efficiency (FUE) and suboptimal yields. This article provides a comprehensive review of the edaphic mechanisms governing P fixation and outlines integrated management strategies. Emphasis is placed on precision placement techniques, soil pH and organic matter optimisation, and the deployment of biological amendments and advanced fertiliser technologies to unlock

"legacy phosphorus" and ensure sustainable agricultural productivity.

Keywords: *Phosphorus fixation, nutrient availability, soil chemistry, fertiliser efficiency, P-solubilising microbes.*

1. Introduction: The Phosphorus Paradox


Phosphorus (P) serves as the biological "energy currency," facilitating energy transfer via adenosine triphosphate (ATP), and it forms the backbone of DNA and RNA. For crops, adequate P supply is non-negotiable for vigorous root establishment, stalk strength, and timely maturation. However, agronomists and producers frequently encounter a confounding scenario known as the "Phosphorus Paradox": soils may possess substantial reserves of total phosphorus, yet crops exhibit classic deficiency symptoms such as stunting and anthocyanin accumulation (purpling).

The root of this inefficiency lies in the soil's chemical environment. Unlike nitrogen, which is highly mobile and prone to

leaching, phosphorus is characterised by low mobility and high reactivity. Upon application, soluble phosphate ions undergo rapid adsorption or precipitation reactions with soil minerals, a process collectively termed **Phosphorus Fixation**. Research indicates that in many mineral soils, crops utilise only 10% to 20% of the applied phosphorus fertiliser in the year of application. The remaining 80% to 90% accumulates as "legacy P"—a banked resource that is chemically locked away. Unlocking this resource is critical not only for farm profitability but also for minimising the environmental footprint associated with excessive fertiliser application.

2. Chemical Mechanisms of Fixation

To effectively manage phosphorus, one must understand the specific chemical pathways that govern its availability. These pathways are strictly dictated by soil pH and mineralogy.



2.1. Acidic Soils (pH < 6.0): The Iron and Aluminum Sink

In acidic environments, the solubility of iron (Fe^{3+}) and aluminium (Al^{3+}) increases significantly. These cations, along with their hydrous oxides (e.g., gibbsite and goethite), possess a strong affinity for the negatively charged orthophosphate ions ($H_2PO_4^-$).

- **Adsorption (Ligand Exchange):**

Phosphate ions displace hydroxyl or water molecules on the surface of clay minerals and iron/aluminium oxides. This forms a stable, inner-sphere complex that is resistant to desorption.

- **Precipitation:** In highly acidic conditions, soluble phosphate reacts directly with dissolved iron and aluminium to precipitate as distinct solid phases, such as variscite ($AlPO_4 \cdot 2H_2O$) and strengite ($FePO_4 \cdot 2H_2O$).

These crystalline compounds have extremely low solubility products, meaning they release P back into the soil solution at negligible rates.

2.2. Alkaline and Calcareous Soils (pH > 7.5): The Calcium Trap

In high-pH soils, calcium becomes the dominant cation. The fixation pathway shifts from adsorption to sequential precipitation.

- **Sequential Precipitation:** When phosphate fertiliser is applied, it initially forms dicalcium phosphate, which is moderately available to plants. However, over time, this metastable compound transforms into tricalcium phosphate and eventually into hydroxyapatite-like minerals. These stable forms mimic raw rock phosphate and are largely unavailable to crops.
- **Surface Adsorption on Carbonates:** In calcareous soils, calcium carbonate (CaCO_3) particles provide vast surface areas where phosphate ions can be adsorbed, further reducing the concentration of P in the soil solution.

2.3. The "Goldilocks Zone" of Availability

Maximum phosphorus availability generally occurs within a neutral pH range of **6.0 to 7.0**. Deviating from this range engages the chemical fixation mechanisms—either the Fe/Al "vice" in acidity or the Ca "lock" in alkalinity.

3. Diagnostic Approaches

Identifying phosphorus fixation requires differentiating between what is present in the soil and what is accessible to the plant.

3.1. Interpreting Soil Analysis

Standard soil tests do not measure total elemental phosphorus; rather, they use specific extraction solutions to estimate the "plant-available" fraction.

- **Bray-P1:** Optimised for acidic to neutral soils; utilises dilute acid and fluoride to solubilise Al-bound P.
- **Olsen-P:** The standard for calcareous soils; utilises a sodium bicarbonate solution to extract P without dissolving calcium carbonates.
- **Mehlich-3:** A robust, multi-element extractant suitable for a wide range of soil types (acidic to neutral).

The Fixation Indicator: A widening gap between the quantity of P fertiliser applied over time, and the slow response of soil test P levels often indicates a soil with a high **Phosphorus Buffering Capacity (PBC)**—essentially, a high capacity to fix added P.

3.2. Symptomatology

Visual diagnosis validates soil data. Since P is mobile within plant tissues, it is translocated from older to newer growth under stress.

- **Anthocyanin Accumulation:** A purpling of older leaves/stems, particularly in maize and brassicas, caused by sugar accumulation resulting from stalled carbohydrate metabolism.

- **Delayed Phenology:** P-deficient crops often exhibit delayed flowering and maturity, increasing the risk of exposure to late-season environmental stresses.

4. Agronomic Management Strategies

Mitigating fixation requires a shift from "feeding the soil" to "feeding the plant."

4.1. Precision Placement: The Banding Advantage

The broadcast application exposes fertiliser granules to maximum soil contact, accelerating fixation. Conversely, **banding** creates a concentrated zone of fertility.

- **Saturation of Fixation Sites:** By concentrating fertiliser in a band (e.g., 5 cm below and to the side of the seed), the adsorption sites on soil minerals within that specific zone become saturated. Excess phosphorus remains in the soil solution, where it is available for uptake.
- **Root Interception:** Banding places nutrients directly in the path of developing roots, reducing the energy cost of nutrient foraging.

4.2. pH Optimization

Correcting soil reaction is the most fundamental method to alter chemical equilibria.

- **Liming:** In acid soils, applying lime neutralises acidity and precipitates

active aluminium, thereby reducing the formation of insoluble Al-phosphates.

- **Rhizosphere Acidification:** In alkaline soils, using ammonium-based fertilisers can lower the pH within the root zone (rhizosphere) during nitrification, temporarily increasing the solubility of calcium-phosphates.

5. The Role of Organic Matter and Biology

Integrating biological principles into chemical management offers a synergistic pathway to P availability.

5.1. Organic Matter and Anion Competition

Soil Organic Matter (SOM) combats fixation through multiple mechanisms:

- **Chelation:** Decomposing organic matter releases organic acids (e.g., humic, fulvic, citric) that form complexes with Fe, Al, and Ca. By "sequestering" these metal ions, organic acids prevent them from reacting with phosphate.
- **Competitive Adsorption:** Organic anions compete with phosphate ions for the same binding sites on clay surfaces, effectively blocking the fixation sites.

5.2. Microbial Solubilization

The soil microbiome plays a pivotal role in P cycling.

- **Arbuscular Mycorrhizal Fungi (AMF):** These symbiotic fungi extend hyphae into soil micropores inaccessible to root hairs, scavenging P and transporting it to the host plant in exchange for carbon.
- **Phosphate-Solubilising Bacteria (PSB):** Genera such as *Bacillus* and *Pseudomonas* secrete organic acids and enzymes (phosphatases) that dissolve mineral phosphates and mineralise organic P, transforming it into plant-available orthophosphate.

6. Advanced Fertiliser Technologies

Innovation in fertiliser formulation aims to physically or chemically protect phosphate ions.

- **Polymer-Coated Fertilisers:** These "controlled-release" granules regulate the diffusion of P into the soil solution, matching nutrient release with crop demand and minimising the time window for fixation reactions.
- **Orthophosphate vs. Polyphosphate:** Liquid polyphosphates contain chains of P molecules. As these chains hydrolyse in soil, they can sequester

cations (Ca, Mg), temporarily protecting P from precipitation.

- **Struvite (Recycled P):** Magnesium ammonium phosphate (struvite), recovered from wastewater, exhibits unique slow-release properties. It is sparingly soluble in water but becomes soluble in the presence of root exudates (citric acid), providing a "demand-driven" nutrient supply.

7. Conclusion

Optimising phosphorus management is essential for closing the yield gap in modern agriculture. The challenge of phosphorus fixation cannot be solved by simply increasing application rates; it requires a strategic, integrated approach. By combining chemical diagnostics (appropriate soil testing), physical precision (banding), biological enhancement (OM and microbial inoculants), and technological innovation (advanced fertilisers), producers can improve Phosphorus Use Efficiency (PUE).

This holistic strategy not only revitalises the productivity of "locked" soils but also ensures the economic and environmental sustainability of farming systems for future generations.

References

1. Fraser, T. D., Lynch, D. H., O'Halloran, I. P., Voroney, R. P.,

- Entz, M. H., & Dunfield, K. E. (2019). Soil phosphorus bioavailability as influenced by long-term management and applied phosphorus source. *Canadian Journal of Soil Science*, 99(3), 292–304. <https://doi.org/10.1139/cjss-2018-0075>
2. Hernández Jiménez, J. E., Nyiraneza, J., Fraser, T. D., Peach Brown, H. C., Lopez-Sanchez, I. J., & Botero-Botero, L. R. (2021). Enhancing phosphorus release from struvite with biostimulants. *Canadian Journal of Soil Science*, 101(1), 22–32. <https://doi.org/10.1139/cjss-2019-0147>
3. Johan, P. D., Ahmed, O. H., Omar, L., & Hasbullah, N. A. (2021). Phosphorus transformation in soils following co-application of charcoal and wood ash. *Agronomy*, 11(10), 2010. <https://doi.org/10.3390/agronomy11102010>
4. Shen, J., Yuan, L., Zhang, J., Li, H., Bai, Z., Chen, X., Zhang, W., & Zhang, F. (2011). Phosphorus dynamics: From soil to plant. *Plant Physiology*, 156(3), 997–1005. <https://doi.org/10.1104/pp.111.175232>
5. Wang, C., Pan, G., Lu, X., & Qi, W. (2023). Phosphorus solubilizing microorganisms: Potential promoters of agricultural and environmental engineering. *Frontiers in Bioengineering and Biotechnology*, 11, 1181078. <https://doi.org/10.3389/fbioe.2023.1181078>