

# AGRIFRONTLINE

*The Voice of Agriculture*

## Beneficial insects: Natural pest control methods to enhance sustainable farming

*Ningaraj Belagalal<sup>1</sup>\*, Maarujodu Sanjana Bhavani<sup>2</sup>, Nishath Anjum<sup>2</sup>, Jampala Akshitha<sup>2</sup>, Gundu Mukunda Munjal<sup>2</sup>*

### Abstract

Beneficial insects play a central role in sustainable agriculture by providing natural pest control, pollination, and ecosystem regulation. Their use reduces dependence on synthetic pesticides, lowers production costs, and limits negative impacts on human health and the environment. This article examines the major groups of beneficial insects, their mechanisms of pest suppression, and their integration into modern farming systems. It discusses ecological principles such as biological control, habitat management, and biodiversity conservation that support beneficial insect populations. Empirical evidence from global cropping systems demonstrates yield stability, pest reduction, and economic benefits. The article also addresses challenges, including climate change, pesticide misuse, and landscape simplification. Practical strategies for farmers and policymakers are outlined to promote adoption at scale. The analysis highlights beneficial insects as a scientifically proven, economically viable, and environmentally sound foundation for sustainable farming worldwide.

**Keywords:** Beneficial insects, Biological control. Sustainable agriculture, Integrated pest management, Agroecosystems

### 1. Introduction:

Modern agriculture faces a persistent challenge. Crop losses due to insect pests remain high despite decades of chemical pesticide use. The Food and Agriculture Organisation estimates that 20 to 40 per cent of global crop production is lost annually to pests. Farmers respond by applying pesticides, often repeatedly. This approach increases costs and creates resistance in pest populations. It also contaminates soil and water, reduces biodiversity, and threatens farm worker health.

Sustainable farming requires alternatives that maintain productivity while reducing environmental harm. Biological processes offer such options. Among them, beneficial insects stand out as a

practical and well-studied solution. These organisms naturally regulate pest populations through predation, parasitism, and competition. They operate continuously and adapt to local ecosystems.

Interest in beneficial insects has grown due to rising input costs, stricter pesticide regulations, and consumer demand for safe food. Advances in ecological science and agronomy now allow farmers to conserve and deploy beneficial insects intentionally. This article examines how beneficial insects function, the evidence supporting their effectiveness, and how they can be scaled across diverse farming systems.

<sup>1</sup> Assistant Professor, Department of Entomology  
KLE College of Agricultural Sciences, Tenikolla, Belagavi

<sup>2</sup> B.Sc (Hons) Agriculture  
School of Agriculture, SR University, Warangal 506371

## Types of beneficial insects and their ecological roles

Beneficial insects include predators, parasitoids, pollinators, and decomposers. Each group contributes differently to agroecosystem stability.

Predatory insects feed directly on pest species. Common examples include lady beetles, lacewings, ground beetles, and predatory bugs. A single lady beetle larva can consume several hundred aphids during its development. Ground beetles prey on soil-dwelling pests such as cutworms and root maggots. These predators are especially important in cereals, vegetables, and orchards.

Parasitoids lay their eggs inside or on pest insects. The developing larvae consume the host, leading to its death. Parasitic wasps and flies are highly host-specific. This specificity reduces the risk of non-target effects. For example, *Trichogramma* wasps are widely used to control lepidopteran pests in maize, cotton, and sugarcane.

Pollinators such as bees, hoverflies, and butterflies indirectly support pest management by improving plant health and yield. Healthy plants tolerate pest damage better. Hoverfly larvae also prey on aphids, providing dual benefits.

Decomposers such as dung beetles and detritivorous insects recycle organic matter. Their activity improves soil structure and nutrient availability. Healthy soils support more vigorous plants and higher populations of natural enemies.

Together, these insects form complex food webs that stabilise agroecosystems. Their presence depends on habitat quality, crop diversity, and management practices.

## Mechanisms of natural pest control

Beneficial insects suppress pest populations through several mechanisms. Understanding these processes helps optimise their use in farming systems.

Predation is the most visible mechanism. Predators reduce pest numbers by consuming them at multiple

life stages. Unlike pesticides, predators respond dynamically to pest density. When pest populations increase, predator reproduction and feeding rates also increase. This density-dependent response prevents pest outbreaks.

Parasitism acts more slowly but provides long-term control. Parasitoids often target specific pest species and life stages. This precision limits pest recovery and reduces the likelihood of resistance. Studies in rice systems show that parasitoids can suppress planthopper populations below economic thresholds without chemical intervention.

Competition and behavioural effects also matter. The presence of natural enemies alters pest behaviour. Pests may reduce feeding or disperse to less favourable habitats—these non-lethal effects reduce crop damage even when pest numbers remain unchanged.

Habitat-mediated control enhances these mechanisms. Flowering plants provide nectar and pollen for adult parasitoids and predators. Hedgerows and field margins offer shelter and overwintering sites. Research in European farms shows that fields with diverse margins host higher densities of beneficial insects and experience lower pest pressure.

## Evidence from global farming systems

Empirical evidence supports the effectiveness of beneficial insects across regions and crops. In Asian rice systems, conserving natural enemies reduced pesticide use by up to 70 per cent while maintaining yields. Farmer field schools in Southeast Asia demonstrated that early-season pesticide sprays disrupted predator populations and increased pest outbreaks. When farmers stopped spraying, natural enemies recovered, and pest damage declined.

In Latin American coffee plantations, shade-grown systems support higher insect diversity. Predatory ants and parasitoids reduce populations of coffee berry borer, a significant pest. Studies report greater

yield stability and reduced chemical inputs in diversified systems compared to sun-grown monocultures.

In North America, biological control of soybean aphid using lady beetles and parasitic wasps has saved farmers millions of dollars annually. Economic analyses estimate that natural enemies provide ecosystem services worth billions of dollars each year in the United States alone.

African smallholder farms also benefit. Intercropping maize with flowering plants increases parasitoid abundance and reduces stem borer damage. This approach improves food security without requiring expensive inputs.

These examples show that beneficial insects are effective across climatic zones, farm sizes, and cropping systems. Success depends on aligning management practices with ecological principles.

### **Integration into integrated pest management programs**

Integrated pest management, or IPM, provides a framework for using beneficial insects effectively. IPM combines biological, cultural, mechanical, and chemical tools, using monitoring and thresholds.

Beneficial insects form the backbone of IPM. Farmers first monitor pest and natural enemy populations. If natural enemies are abundant and pest levels remain below economic thresholds, no action is needed. This approach avoids unnecessary interventions.

Cultural practices support beneficial insects. Crop rotation breaks pest life cycles. Intercropping increases habitat diversity. Reduced tillage preserves ground-dwelling predators. Selecting pest-resistant crop varieties lowers pest pressure and supports natural control.

Chemical pesticides remain a tool, but are used selectively. When needed, farmers choose products with low toxicity to natural enemies. Application timing avoids the peak activity of beneficial insects.

This selective use preserves biological control agents.

Augmentative biological control involves releasing commercially reared beneficial insects. This method is standard for greenhouse and high-value crops. While effective, it requires technical knowledge and can be costly. Conservation biological control, which focuses on protecting existing populations, is often more sustainable for open field systems.

Successful IPM programs rely on farmer education and extension services. Training improves pest identification and decision-making. Countries that invested in IPM training have reduced pesticide use while maintaining productivity.

### **Constraints and emerging challenges**

Despite clear benefits, the adoption of beneficial insect-based pest control faces constraints. One major challenge is pesticide dependency. Farmers accustomed to quick chemical solutions may perceive biological control as slow or uncertain. This perception persists even when evidence shows long-term benefits.

Landscape simplification reduces habitat for beneficial insects. Large monocultures lack floral resources and refuges. Restoring habitat requires coordinated action beyond individual farms. Policy incentives are often necessary.

Climate change introduces uncertainty. Rising temperatures alter insect life cycles and distribution. Some pests expand their range faster than their natural enemies. Research is needed to understand these dynamics and adapt management strategies.

Knowledge gaps also limit adoption. Identifying beneficial insects requires training. Misidentification can lead to unnecessary spraying. Extension systems in many regions remain underfunded.

Finally, market structures influence farmer decisions. Short-term yield maximisation often outweighs long-term sustainability. Aligning

economic incentives with ecological outcomes remains a policy challenge.

## Conclusion

Beneficial insects offer a scientifically grounded solution to one of agriculture's most persistent problems. They provide effective pest control, reduce reliance on synthetic pesticides, and support broader ecosystem services. Evidence from diverse farming systems shows that when managed properly, beneficial insects enhance yield stability and farm profitability.

Their success depends on ecological understanding and supportive management practices. Habitat conservation, crop diversity, and selective pesticide use are essential. Integrated pest management provides a practical framework for implementation.

## References

1. Altieri, M. A., & Nicholls, C. I. (2020). Agroecology and the emergence of a post-COVID agriculture. *Agriculture and Human Values*, 37(3), 525–526.
2. Bale, J. S., van Lenteren, J. C., & Bigler, F. (2008). Biological control and sustainable food production. *Philosophical Transactions of the Royal Society B*, 363(1492), 761–776.
3. Costanza, R., et al. (2014). Changes in the global value of ecosystem services. *Global Environmental Change*, 26, 152–158.
4. Crowder, D. W., & Jabbour, R. (2014). Relationships between biodiversity and biological control in agroecosystems. *Current Opinion in Insect Science*, 2, 1–7.
5. FAO. (2019). Integrated pest management and pesticide risk reduction. Food and Agriculture Organisation of the United Nations.
6. Gurr, G. M., Wratten, S. D., Landis, D. A., & You, M. (2017). Habitat management to suppress pest populations. *Annual Review of Entomology*, 62, 91–109.
7. Heimpel, G. E., & Mills, N. J. (2017). *Biological control: Ecology and applications*. Cambridge University Press.
8. Kremen, C., & Miles, A. (2012). Ecosystem services in biologically diversified farming systems. *Ecology and Society*, 17(4), 40.
9. Landis, D. A., Wratten, S. D., & Gurr, G. M. (2000). Habitat management to conserve natural enemies of arthropod pests. *Annual Review of Entomology*, 45, 175–201.
10. Losey, J. E., & Vaughan, M. (2006). The economic value of ecological services provided by insects. *BioScience*, 56(4), 311–323.
11. Naranjo, S. E., Ellsworth, P. C., & Frisvold, G. B. (2015). Economic value of biological control in integrated pest management of managed plant systems. *Annual Review of Entomology*, 60, 621–645.

Challenges remain, including climate change, landscape simplification, and limited extension capacity. Addressing these issues requires coordinated efforts among farmers, researchers, policymakers, and markets. Investment in education and incentives can accelerate adoption.

Sustainable farming is not a return to low productivity systems. It is a transition toward knowledge-intensive agriculture that works with biological processes. Beneficial insects are central to this transition. By recognising and supporting their role, agriculture can move toward resilience, environmental integrity, and long-term food security.